

# Proinflammatory cytokines, sickness behavior, and Alzheimer disease

C. Holmes, MRCPsych,  
PhD  
C. Cunningham, PhD  
E. Zotova, BSc  
D. Culliford, MSc  
V.H. Perry, DPhil

Address correspondence and reprint requests to Prof. Clive Holmes, University of Southampton, Memory Assessment and Research Centre, Botley Rd, Southampton, UK, SO30 3JB  
ch4@soton.ac.uk

## ABSTRACT

**Background:** In Alzheimer disease (AD), systemic inflammation is known to give rise to a delirium. However, systemic inflammation also gives rise to other centrally mediated symptoms in the absence of a delirium, a concept known as sickness behavior. Systemic inflammation is characterized by the systemic production of the proinflammatory cytokines tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) and interleukin-6 (IL-6) that mediate immune to brain communication and the development of sickness behavior.

**Objective:** To determine if raised serum TNF $\alpha$  or IL-6 are associated with the presence of sickness behavior symptoms, independent of the development of delirium, in a prospective cohort study of subjects with AD.

**Methods:** A total of 300 subjects with mild to severe AD were cognitively assessed at baseline and a blood sample taken for inflammatory markers. Cognitive assessments, including assessments to detect the development of a delirium, and blood samples were repeated at 2, 4, and 6 months. The development of neuropsychiatric symptoms in the subject with AD over the 6-month follow-up period was assessed independently by carer interview at 2, 4, and 6 months.

**Results:** Raised serum TNF $\alpha$  and IL-6, but not CRP, were associated with an approximately 2-fold increased frequency of neuropsychiatric symptoms characteristic of sickness behavior. These relationships are independent of the development of delirium.

**Conclusions:** Increased serum proinflammatory cytokines are associated with the presence of symptoms characteristic of sickness behavior, which are common neuropsychiatric features found in AD. This association was independent of the presence of delirium. *Neurology*® 2011;77:212-218

## GLOSSARY

**AD** = Alzheimer disease; **ADAS-Cog** = Alzheimer's Disease Assessment Scale Cognitive subscale; **CAM** = Confusion Assessment Method; **Chel** = cholinesterase inhibitor; **CI** = confidence interval; **CRP** = C-reactive protein; **DSM-III-R** = *Diagnostic and Statistical Manual of Mental Disorders*, 3rd edition, revised; **DSM-IV** = *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition; **IL-6** = interleukin-6; **IQR** = interquartile range; **MSD** = Meso Scale Discovery; **NINCDS-ADRDA** = Neurological and Communicative Disorders and Stroke-Alzheimer's Disease and Related Disorders Association; **NPI** = Neuropsychiatric Inventory; **OR** = odds ratio; **SIE** = systemic inflammatory event; **TNF $\alpha$**  = tumor necrosis factor- $\alpha$ .

Systemic inflammation is characterized by the production of C-reactive protein (CRP) and the proinflammatory cytokines tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) and interleukin 6 (IL-6). TNF $\alpha$  and IL-6 play a role in immune to brain communication by activating the central innate immune response to initiate a behavioral response known as sickness behavior.<sup>1,2</sup> Sickness behavior refers to a coordinated set of behavioral changes that develop during the course of raised systemic inflammation.<sup>3</sup> These behaviors include increased anxiety, depressed mood, and apathy, and are adaptive protective mechanisms aimed at conserving energy and reducing further exposure to systemic inflammatory insults<sup>4</sup> (table e-1 on the *Neurology*® Web site at www.neurology.org). In animal models of neurodegeneration, systemic inflammation results in a markedly increased central proinflammatory cytokine profile and the development of sickness

Editorial, page 206

See pages 219 and 227

Supplemental data at www.neurology.org



Scan this code with your smartphone to access this feature

From the Clinical Neurosciences Division (C.H., E.Z.) and School of Biological Science (V.H.P.), University of Southampton, Southampton; Memory Assessment and Research Centre (C.H.), Moorgreen Hospital, Hampshire Partnership Foundation Trust, Southampton; Trinity College Institute of Neuroscience (C.C.), School of Biochemistry & Immunology, Trinity College Dublin, Dublin; and Research Design Service (D.C.), Southampton General Hospital, Southampton, UK.

**Study funding:** Supported by the UK Alzheimer's Society.

**Disclosure:** Author disclosures are provided at the end of the article.

behavior and neuronal cell loss.<sup>5,6</sup> In Alzheimer disease (AD), we have shown that systemic inflammation, associated with raised serum proinflammatory cytokines, is associated with a marked increase in cognitive decline that is independent of acute cognitive deterioration associated with delirium.<sup>7,8</sup> We hypothesized that in AD raised serum TNF $\alpha$  and IL-6 would also be associated with an exacerbation of neuropsychiatric symptoms characteristic of sickness behavior independent of the development of the behavioral symptoms of delirium.

**METHODS Study design.** A total of 300 community-dwelling subjects with mild to severe dementia and their caregivers were recruited between November 2003 and May 2006 from clinical referrals to memory assessment services in Southampton, UK. Following consent procedures, all subjects fulfilling National Institute of Neurological and Communicative Disorders and Stroke–Alzheimer’s Disease and Related Disorders Association (NINCDS-ADRDA)<sup>9</sup> criteria for probable or possible AD were tested using the Alzheimer’s Disease Assessment Scale Cognitive subscale (ADAS-Cog).<sup>10</sup> Immediately following cognitive assessment, a blood sample for assaying CRP and the proinflammatory cytokines TNF $\alpha$  and IL-6 was taken. At the end of the interview, the main caregiver was given a once-weekly checklist diary based on the Neuropsychiatric Inventory (NPI)<sup>11</sup> and the Confusion Assessment Method (CAM)<sup>12</sup> (see table e-1) in which to enter the presence of behavioral symptoms over the following 2 months and to alert the study coordinator if a delirium was suspected. The subjects’ main caregiver was then formally interviewed 2 months later by a separate research nurse, blind to the initial cognitive assessment of the subject, using the NPI to determine (with cross-reference to the checklist) the presence of neuropsychiatric features and acute systemic inflammatory events (SIEs) over the preceding 2 months. An SIE was defined as a short-lived (less than 2 months duration) infection or trauma not directly involving the CNS with a minimum serum CRP level of 1  $\mu$ g/mL after the event. The NPI is a carer-based interview that assesses 10 behavioral disturbances in the subject. Based on animal and human clinical studies,<sup>13–17</sup> we identified 3 core symptoms from the NPI as being a priori compatible with the concept of sickness behavior (i.e., increased depression/dysphoria, increased anxiety, and increased apathy). The main caregiver was revisited at 4 and 6 months with the NPI reassessed in an identical manner to that at 2 months.

All patients were cognitively reassessed 2, 4, and 6 months after baseline (within 2 days and independently of the NPI carer assessment) by the original patient assessor using the ADAS-Cog and the CAM to assess of the development of a delirium with additional blood sampling for inflammatory markers. The CAM operationalizes the core symptoms of delirium as defined by *DSM-III-R*<sup>18</sup> and has a high concordance with the *DSM-IV* diagnosis of delirium.<sup>19–21</sup> Additional assessments using the CAM were carried out if a delirium was suspected by the main carer in between scheduled visits.

**Standard protocol approvals, registrations, and patient consents.** This study received approval from the South and West Hampshire Local Research Ethics Committee (ref 237/03/w).

Written informed consent was obtained from all patients (or guardians of patients) participating in the study.

**Systemic inflammation assays.** Blood sera were immediately placed on ice and stored within 2 hours at  $-80^{\circ}\text{C}$ . CRP was assayed using ELISA and had a detection limit of 1  $\mu$ g/mL. TNF $\alpha$  and IL-6 was assayed using sandwich immunoassay multiplex cytokine assay (Meso Scale Discovery [MSD]). A protocol provided by MSD for custom assays was used with no major modifications. The lowest detectable limit was 1.1 pg/mL for TNF $\alpha$  and 0.7 pg/mL for IL-6.

**Statistical analysis.** Normality of continuous variables was determined by quantile-quantile plots of the residuals. Baseline age and ADAS-Cog were normally distributed. Serum CRP, TNF $\alpha$ , and IL-6 levels were not normally distributed. Following previous guidelines,<sup>22</sup> low serum CRP was defined as  $<1.0$   $\mu$ g/mL at all 4 timepoints; moderate/high levels were defined as above 1.0  $\mu$ g/mL at any timepoint. Serum TNF $\alpha$  and IL-6 levels were simplified by the use of quartile ranges, based on subject numbers found at baseline assessment. Low serum TNF $\alpha$  was defined as that found in the lowest quartile for TNF $\alpha$  ( $<2.4$  pg/mL) at all 4 timepoints and high serum levels as above 2.4 pg/mL at any timepoint. Low serum IL-6 was defined as that found in the lowest quartile for IL-6 ( $<2.8$  pg/mL) at all 4 timepoints and high serum levels as above 2.8 pg/mL at any timepoint. The average NPI score for each individual over the 6-month follow-up period was determined by taking the mean of the 3 NPI scores (product of the frequency and severity) at the 3 timepoints at 2, 4, and 6 months and was not normally distributed. Impairment of an individual NPI symptom was based on its presence (i.e., frequency of  $\geq 1$ ) at any time during the 6-month follow-up period. Allowing for a 10% dropout rate, 300 subjects gave 90% power to detect a significant difference ( $\alpha = 0.017$ ) of 15% between the total NPI score in subjects in the lowest quartile compared to subjects in the top 3 quartiles. Relationships between variables was assessed using a mixture of Student  $t$  test, Fisher exact test,  $\chi^2$ , Mann Whitney  $U$ , and Spearman correlations. Data interaction was assessed using logistic regression analysis.

**RESULTS** A total of 366 subjects were approached for inclusion. A total of 51 subjects or their carers declined participation; 15 subjects did not fulfill NINCDS-ADRDA criteria. The 51 nonparticipating subjects did not differ with respect to age (nonparticipants age 82.6 [SE 0.9] years vs participants 82.8 [SE 0.4]; mean difference 0.2 [95% confidence interval (CI)  $-2.1$  to 2.3 years],  $t$  test  $p = 0.9$ ) or gender (female nonparticipants 38/51 [75%] vs female participants 198/300 [67%],  $\chi^2$  1.4,  $p = 0.2$ ). A total of 25 subjects were clinically unresponsive at baseline with severe end-stage dementia, i.e., ADAS-Cog scores greater than 60 points, and were excluded from further analysis due to the difficulty in assessing the development of a delirium over the follow-up period.

**Baseline data.** Of the 275 subjects, 161 (59%) fulfilled NINCDS-ADRDA criteria for probable AD and 114 (41%) possible AD. The mean age of the cohort at baseline was 82.7 (SD 7.4) years. A total of

**Table 1** Baseline serum systemic inflammatory markers and demographics

Systemic levels of inflammatory marker (n)	Age (SE), y	Sex, n (%) women	Baseline ADAS-Cog score (SE), patients
<b>CRP</b>			
Low <1 µg/mL (83)	81.0 (0.8) <sup>a</sup>	50 (60.2) <sup>d</sup>	28.0 (1.5) <sup>g</sup>
High ≥1 µg/mL (186)	83.5 (0.6)	124 (66.7)	30.7 (0.9)
<b>TNFα</b>			
Low <2.4 pg/mL (65)	79.5 (1.0) <sup>b</sup>	43 (66.2) <sup>e</sup>	25.6 (1.6) <sup>h</sup>
High ≥2.4 pg/mL (204)	83.9 (0.5)	131 (64.2)	31.0 (0.9)
<b>IL-6</b>			
Low <2.8 pg/mL (66)	81.1 (0.9) <sup>c</sup>	43 (65.2) <sup>f</sup>	30.0 (1.5) <sup>i</sup>
High ≥2.8 pg/mL (203)	83.3 (0.5)	131 (64.5)	29.3 (0.9)

Abbreviations: ADAS-Cog = Alzheimer's Disease Assessment Scale Cognitive subscale; CI = confidence interval; CRP = C-reactive protein; IL-6 = interleukin-6; TNFα = tumor necrosis factor α.

<sup>a</sup> Mean difference 2.5 (95% CI 0.6 to 4.5) years; *t* test *p* = 0.01.

<sup>b</sup> Mean difference 4.4 (95% CI 2.4 to 6.4) years; *t* test *p* < 0.0001.

<sup>c</sup> Mean difference 2.2 (95% CI 0.2 to 4.3) years; *t* test *p* = 0.03.

<sup>d</sup>  $\chi^2$  1.0, *p* = 0.3.

<sup>e</sup>  $\chi^2$  0.08, *p* = 0.8.

<sup>f</sup>  $\chi^2$  0.01, *p* = 0.9.

<sup>g</sup> Mean difference 2.7 (95% CI -6.1 to 0.7) patients; *t* test *p* = 0.1.

<sup>h</sup> Mean difference 5.4 (95% CI 1.9 to 9.0) patients; *t* test *p* = 0.003.

<sup>i</sup> Mean difference -0.7 (95% CI -2.9 to 1.5) patients; *t* test *p* = 0.7.

99 (36%) subjects were men. At baseline, subjects had a mean ADAS-Cog score of 29.6 (SD 13.0) points. Blood sampling was obtained in 269/275 (98%) subjects at baseline. Median serum CRP was 2.5 (interquartile range [IQR] 0–5.8) µg/mL and was moderate/high (serum level ≥1 µg/mL) in 186 (70%) subjects. Serum TNFα and IL-6 was detectable in all subjects (TNFα median 3.3 pg/mL [IQR 2.4 pg/mL–4.2 pg/mL]; IL-6 median 4.5 pg/mL [IQR 2.8 pg/mL–8.0 pg/mL]).

Low levels of all systemic inflammatory markers at baseline were associated with a younger age, but not gender. Low baseline levels of TNFα, but not

**Table 2** Frequency of neuropsychiatric features in subjects with or without presence of delirium

Neuropsychiatric feature	Delirium absent (n = 197), n (%)	Delirium present (n = 25), n (%)	$\chi^2$ , <i>p</i>
Delusions	77 (39)	15 (60)	$\chi^2$ 4.0, <i>p</i> = 0.046
Hallucinations	62 (32)	13 (52)	$\chi^2$ 4.2, <i>p</i> = 0.04
Agitation	98 (50)	21 (84)	$\chi^2$ 10.5, <i>p</i> = 0.001
Depression	94 (48)	18 (72)	$\chi^2$ 5.2, <i>p</i> = 0.02
Anxiety	108 (55)	20 (80)	$\chi^2$ 5.8, <i>p</i> = 0.02
Elation	35 (18)	6 (24)	$\chi^2$ 0.6, <i>p</i> = 0.5
Apathy	116 (59)	16 (64)	$\chi^2$ 0.2, <i>p</i> = 0.6
Disinhibition	44 (22)	13 (52)	$\chi^2$ 10.2, <i>p</i> = 0.001
Irritability	86 (44)	17 (68)	$\chi^2$ 5.3, <i>p</i> = 0.02
Motor activity	81 (41)	19 (76)	$\chi^2$ 10.9, <i>p</i> = 0.001

CRP or IL-6, were associated with lower ADAS-Cog scores at baseline (table 1).


**Six-month follow-up period.** A total of 222 (81%) subjects had complete clinical and systemic inflammatory marker follow-up at 2, 4, and 6 months. Of the 53 subjects who did not complete the study, 15 subjects died before study completion and 38 subjects refused phlebotomy or further cognitive assessment at some point during the study. There was no significant difference between those subjects who completed the study in terms of age (completers 82.4 [SE 0.5] years vs noncompleters 83.9 [SE 1.0] years; mean difference 1.5 [95% CI -0.7 to 3.7] years, *t* test *p* = 0.2), gender (female completers 142 [64%] vs female noncompleters 34 [64%],  $\chi^2$  0.001, *p* = 1.0), or baseline ADAS-Cog score (completers 28.9 [SE 0.8] patients vs noncompleters 32.2 [SE 2.0] patients; mean difference 3.3 [95% CI -0.6 to 7.2] patients, *t* test *p* = 0.1).

Of the 222 subjects who completed the study, 110 (45.5%) had a total of 150 systemic inflammatory events; 90 (41%) were taking a cholinesterase inhibitor (ChEI) and 88 (40%) an antidepressant. Sixty-six subjects (30%) had a history of hypertension, 35 (16%) hypercholesterolemia, and 16 (7%) type 2 diabetes. The use of ChEI or antidepressants was not related to the presence of the 3 core features of sickness behavior (symptoms present in 78 [87%] subjects taking a ChEI vs 105 [80%] subjects not taking a ChEI,  $\chi^2$  1.9, *p* = 0.2; symptoms present in 70 [80%] subjects taking an antidepressant vs 113 [84%] subjects not taking an antidepressant,  $\chi^2$  0.8, *p* = 0.4).

**Neuropsychiatric features and delirium.** The median NPI score over the 6-month period was 8.2 (IQR 3.3–13.7) patients and was significantly correlated with age at baseline (Spearman correlation -0.18, *p* = 0.007). Delirium was identified in 25 (11.2%) subjects during the 6-month follow-up period. No relationship was found with age and development of delirium (delirium absent age 83.0 [SE 0.5] years vs delirium present 81.0 [SE 1.6] years; mean difference 1.6 years [95% CI -1.5 to 4.7], *p* = 0.3).

Subjects with delirium had a higher median NPI score over the 6-month follow-up period than those without identified delirium (delirium absent NPI score 6.7 [IQR 2.7–12.7] patients vs delirium present 11.0 [IQR 15.7–22.7] patients, MWU *p* < 0.0001). The frequency of individual symptoms in subjects with or without delirium is shown in table 2 and figure e-1. All symptoms were numerically more frequent in subjects with delirium compared to those without delirium and this was

**Figure 1** Frequency distribution of neuropsychiatric features in subjects with Alzheimer disease over the 6-month follow-up period by low or high levels of tumor necrosis factor- $\alpha$  (TNF $\alpha$ )



\* $p \leq 0.05$ , \*\* $p \leq 0.01$ , adjusted for baseline age, gender, Alzheimer's Disease Assessment Scale Cognitive subscale score, and presence of delirium during follow-up.

statistically significant for all symptoms except for elation and apathy. The 3 core features of sickness behavior were more common in subjects with an identified delirium over the 6-month follow-up period than those without an identified delirium (sickness behavior symptoms present in 24 [96%] subjects with identified delirium compared to 159 [81%] subjects without an identified delirium; Fisher exact test  $p = 0.04$ ). There was a significant relationship between baseline ADAS-Cog score and NPI score during the 6-month follow-up period (Spearman correlation 0.24,  $p < 0.0001$ ) and between baseline ADAS-Cog score and the development of a delirium (delirium present baseline ADAS-Cog score 36.4 patients vs delirium absent

28.0 patients [mean difference 8.4 points (3.3 to 13.6)],  $p = 0.001$ ).

**Systemic inflammatory markers.** During the 6-month follow-up period, 27 subjects with low serum CRP levels at baseline had serum CRP levels that remained at less than 1.0  $\mu\text{g/mL}$ , 34 subjects with low serum TNF $\alpha$  levels at baseline had serum TNF $\alpha$  levels that remained at less than 2.4  $\text{pg/mL}$ , and 29 subjects with low serum IL-6 levels at baseline had serum IL-6 levels that remained at less than 2.8  $\text{pg/mL}$ .


**Systemic inflammatory markers and delirium.** Subjects with low CRP throughout the study were not less likely to experience an episode of delirium than subjects with high CRP (low CRP 3.7% vs high CRP 12.6%,  $\chi^2$  1.8,  $p = 0.20$ ). Likewise, subjects with low IL-6 throughout the study were not less likely to experience an episode of delirium than subjects with high IL-6 (low IL-6 3.4% vs high IL-6 12.4%,  $\chi^2$  2.0,  $p = 0.15$ ). There was a trend to suggest that subjects with low TNF $\alpha$  throughout the study were less likely to experience an episode of delirium than subjects with high TNF $\alpha$  (low TNF $\alpha$  2.9% vs high TNF $\alpha$  12.8%,  $\chi^2$  2.8,  $p = 0.09$ ).

**Systemic inflammatory markers and neuropsychiatric features.** There was no statistical difference in the NPI score in subjects with low CRP levels throughout the 6-month follow-up period compared to subjects with high CRP levels (low CRP 6.0 [IQR 2.0–14.0] vs high CRP 8.3 [IQR 3.3–13.7], MWU  $p = 0.5$ ).

Subjects with low TNF $\alpha$  levels throughout the 6-month follow-up period had a lower NPI score over this same time period (low TNF $\alpha$  4.8 [IQR 1.1–9.4] vs high TNF $\alpha$  8.8 [IQR 3.7–14.0], MWU  $p = 0.01$ ). The frequency of individual symptoms in subjects with low compared to high levels of TNF $\alpha$  is shown in figure 1. With the exception of disinhibition, all other symptoms were numerically more frequent in subjects with high levels of TNF $\alpha$  compared to those with low levels and this was statistically significant for the presence of agitation ( $\chi^2$  5.4,  $p = 0.02$ ), depression/dysphoria ( $\chi^2$  7.1,  $p = 0.008$ ), and anxiety ( $\chi^2$  6.2,  $p = 0.01$ ) with a trend for apathy ( $\chi^2$  2.6,  $p = 0.1$ ). The relationships between the presence of these symptoms and high TNF $\alpha$  levels was largely unchanged following adjustment for baseline ADAS-Cog score, age, gender, and the presence of delirium during the 6-month follow-up period: agitation (odds ratio [OR] 2.6 [95% CI 1.1 to 5.8],  $p = 0.02$ ), depression/dysphoria (OR 3.1 [95% CI 1.3 to 7.0],  $p = 0.008$ ), anxiety (OR 3.0 [95% CI 1.3 to 6.9],  $p = 0.007$ ), and apathy (OR 2.1 [95% CI 1.0 to 4.6],  $p = 0.06$ ).



**Figure 2** Frequency distribution of neuropsychiatric features in subjects with Alzheimer disease over the 6-month follow-up period by low or high levels of interleukin-6 (IL-6)



\* $p \leq 0.05$ , \*\* $p \leq 0.01$ , adjusted for baseline age and gender.

Subjects with low IL-6 levels throughout the 6-month follow-up period also had a lower NPI score over this same time period (low IL-6 4.3 [IQR 0.7–9.2] vs high IL-6 8.7 [IQR 3.7–13.8], MWU  $p = 0.02$ ). The frequency of individual symptoms in subjects with low compared to high levels of IL-6 is shown in figure 2. All symptoms, with the exception of motor activity, were numerically more frequent in subjects with high levels of IL-6 compared to those with low levels, and this was statistically significant for the presence of hallucinations ( $\chi^2 6.0$ ,  $p = 0.015$ ) and apathy ( $\chi^2 4.5$ ,  $p = 0.03$ ) with a trend for delusions ( $\chi^2 2.6$ ,  $p = 0.1$ ). The relationships between the presence of these symptoms and high IL-6 levels were largely unchanged following adjustment for

**Table 3** Period prevalence of systemic inflammatory events during the 6-month follow-up period and number (%) of subjects with this event experiencing one or more of the 3 core features of sickness behavior

Systemic inflammatory event (no. of subjects identified with event)	No. (% of subjects identified with event) of subjects with one or more core features of sickness behavior
No event (112)	87 (78)
Respiratory infection (51)	44 (86)
Genitourinary infection (27)	22 (82)
Accidental trauma (32)	27 (84)
Gastrointestinal infection (14)	14 (100)
Other infections (13)	13 (100)
Surgical intervention (11)	10 (91)
Myocardial infarction (2)	1 (50)
History of hypertension (66)	54 (82)
History of hypercholesterolemia (35)	30 (86)
History of type II diabetes (16)	15 (94)

baseline age and gender (hallucinations, OR 4.0 [95% CI 1.3 to 12.1],  $p = 0.015$ ; apathy, OR 2.9 [95% CI 1.3 to 6.6],  $p = 0.01$ ) and delusions (OR 2.2 [95% CI 0.9 to 5.4],  $p = 0.08$ ).

The period prevalence of the 3 core features of sickness behavior for subjects experiencing specific SIEs is shown in table 3.

**DISCUSSION** The history of cytokine-induced sickness behavior has been extensively reviewed elsewhere.<sup>4,23</sup> We have argued that these symptoms are likely to be exaggerated and prolonged in subjects with AD due to prior microglial activation in AD<sup>24</sup> and to the effects of SIEs and chronic comorbid conditions associated with raised systemic inflammation, e.g., diabetes.<sup>25</sup> We have previously shown that raised serum TNF $\alpha$ , but not CRP, is associated with increased cognitive decline in AD, independent of delirium<sup>8</sup>; elsewhere there is evidence that raised serum TNF $\alpha$  and IL-6, but not CRP, is associated with cognitive impairment in diabetes.<sup>26</sup> Here we show that raised serum TNF $\alpha$  and IL-6, but not CRP, is associated with an approximately 2-fold increase of NPI scores and an increased frequency of neuropsychiatric symptoms characteristic of sickness behavior, independent of delirium. The lack of relationship between raised CRP as an indicator of systemic inflammation and behavioral symptoms may reflect findings in the elderly that CRP is neither a good predictor of systemic inflammation<sup>27,28</sup> or involved in peripheral to brain communication of sickness behavior.

While every effort was made to identify episodes of delirium in this prospective study using well-validated tools, it is possible that some episodes of delirium may have been missed. However, the frequency of delirium found in this study is in keeping with other community studies of AD,<sup>29,30</sup> and it is notable that the core features of sickness behavior symptoms (apathy, anxiety, and depression) were also common (occurring in around 80%) in those subjects without detectable delirium.

As hypothesized, we found that the 3 core symptoms of sickness behavior were more frequent in subjects with AD with a raised proinflammatory cytokine profile. There appears to be some variability in the effects of different inflammatory events on the development of sickness behavior but small numbers preclude any meaningful comparisons. However, high levels of TNF $\alpha$  and IL-6 were also associated with a general increased frequency of a wide range of symptoms which are not commonly identified as characteristic of sickness behavior (figure 2). These symptoms are common in AD, even in the absence of delirium, and may reflect an exaggerated cytokine response in the brain due to priming of the microglia and enhanced sensitivity to modest systemic inflammatory signals.<sup>5,25</sup> The higher co-occurrence of sickness behavior in subjects with delirium raises the possibility that delirium may represent an extreme nonadaptive presentation of sickness behavior. Thus, there may be a continuum of behavioral changes: in healthy young individuals systemic inflammation inducing benign and transient changes in the CNS while similar systemic changes induce more severe symptoms in a population with dementia. Likewise, delirium may require a more severe systemic inflammation in healthy young individuals but only a mild event in individuals with dementia.<sup>31</sup> It is notable that the symptom profile shown by raised TNF $\alpha$  and IL-6 while both exhibiting apathy also differ. Thus, raised TNF $\alpha$  is associated with symptoms more characteristic of depressive symptomatology while raised IL-6 is more characteristic of a psychotic profile (figures 1 and 2) and is consistent with cluster analyses of the NPI in subjects with AD.<sup>32</sup> In addition, subjects with mild cognitive impairment show a NPI profile that is remarkably similar to that associated with sickness behavior,<sup>33</sup> suggesting that the early presentation of these symptoms might represent the neurochemical consequences of microglial activation and consequently explain their poor cognitive outcome.<sup>34</sup>

#### AUTHOR CONTRIBUTIONS

Prof. Holmes: drafting/ revising the manuscript, study concept or design, analysis or interpretation of data, acquisition of data, statistical analysis, study supervision, obtaining funding. Dr. Cunningham: drafting/ revising

the manuscript, study concept or design, contribution of vital reagents/ tools/ patients. E. Zotova: analysis or interpretation of data, acquisition of data. D. Culliford: drafting/ revising the manuscript, analysis or interpretation of data, statistical analysis. Prof. Perry: drafting/ revising the manuscript, study concept or design, analysis or interpretation of data.

#### ACKNOWLEDGMENT

The authors thank the carers, patients, and care staff participating in the study, and research nurses Jan Woolford, Sara Kerr, and Christine Dean for the collection of the clinical data.

#### DISCLOSURE

Prof. Holmes has received research support from the Alzheimer's Society, the UK Department of Health, and the Medical Research Council. Dr. Cunningham receives research support from the Wellcome Trust and his sibling is an employee of Wyeth. E. Zotova and D. Culliford report no disclosures. Prof. Perry has received honoraria from the Danish Research Council and UCB and receives research support from the European Union, the Medical Research Council, the Biotechnology and Biological Sciences Research Council, and Wellcome Trust.

Received July 23, 2010. Accepted in final form December 22, 2010.

#### REFERENCES

1. Kent S, Bluth RM, Kelley KW, Dantzer R. Sickness behavior as a new target for drug development. *Trends Pharmacol Sci* 1992;13:24–28.
2. Dantzer R. Cytokine, sickness behavior, and depression. *Immunol Allergy Clin North Am* 2009;29:247–264.
3. Hart BL. Biological basis of the behavior of sick animals. *Neurosci Biobehav Rev* 1988;12:123–137.
4. Dantzer R, Kelley KW. Twenty years of research on cytokine-induced sickness behavior. *Brain Behav Immun* 2007;21:153–160.
5. Cunningham C, Wilcockson DC, Champion S, Lunn K, Perry VH. Central and systemic endotoxin challenges exacerbate the local inflammatory response and increase neuronal death during chronic neurodegeneration. *J Neurosci* 2005;25:9275–9284.
6. Cunningham C, Champion S, Lunn K, et al. Systemic inflammation induces acute behavioral and cognitive changes and accelerates neurodegenerative disease. *Biol Psychiatry* 2009;65:304–312.
7. Holmes C, El-Okil M, Williams AL, Cunningham C, Wilcockson D, Perry VH. Systemic infection, interleukin 1beta, and cognitive decline in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2003;74:788–789.
8. Holmes C, Cunningham C, Zotova E, et al. Systemic inflammation and disease progression in Alzheimer disease. *Neurology* 2009;73:768–774.
9. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;34:939–944.
10. Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. *Am J Psychiatry* 1984;141:1356–1364.
11. Cummings JL, Mega M, Gray K, Rosenberg-Thompson S, Carusi DA, Gornbein J. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology* 1994;44:2308–2314.
12. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: the confusion assess-

- ment method: a new method for detection of delirium. *Ann Intern Med* 1990;113:941–948.
13. Piser TM. Linking the cytokine and neurocircuitry hypotheses of depression: a translational framework for discovery and development of novel anti-depressants. *Brain Behav Immun* 2010;24:515–524.
  14. Capuron L, Gumnick JF, Musselman DL, et al. Neurobehavioral effects of interferon-alpha in cancer patients: phenomenology and paroxetine responsiveness of symptom dimensions. *Neuropsychopharmacology* 2002;26:643–652.
  15. Capuron L, Miller AH. Cytokines and psychopathology: lessons from interferon-alpha. *Biol Psychiatry* 2004;56:819–824.
  16. Johnson DR, O'Connor JC, Dantzer R, Freund GG. Inhibition of vagally mediated immune-to-brain signaling by vanadyl sulfate speeds recovery from sickness. *Proc Natl Acad Sci USA* 2005;102:15184–15189.
  17. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci* 2008;9:46–56.
  18. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-III-R*. Washington, DC: American Psychiatric Association; 1987.
  19. Laurila JV, Pitkala KH, Strandberg TE, Tilvis RS. Confusion assessment method in the diagnostics of delirium among aged hospital patients: would it serve better in screening than as a diagnostic instrument? *Int J Geriatr Psychiatry* 2002;17:1112–1119.
  20. Monette J, Galbaud du Fort G, Fung SH, et al. Evaluation of the Confusion Assessment Method (CAM) as a screening tool for delirium in the emergency room. *Gen Hosp Psychiatry* 2001;23:20–25.
  21. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV*. Washington, DC: American Psychiatric Association; 1994.
  22. Pearson TA, Mensah GA, Alexander RW, et al. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: a statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 2003;107:499–511.
  23. Dantzer R. Cytokine-induced sickness behavior: mechanisms and implications. *Ann NY Acad Sci* 2001;933:222–234.
  24. Akiyama H, Barger S, Barnum S et al. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000;21:383–421.
  25. Perry VH, Cunningham C, Holmes C. Systemic infections and inflammation affect chronic neurodegeneration. *Nat Rev Immunol* 2007;7:161–167.
  26. Marioni RE, Strachan MW, Reynolds RM, et al. Association between raised inflammatory markers and cognitive decline in elderly people with type 2 diabetes: the Edinburgh Type 2 Diabetes Study. *Diabetes* 2010;59:710–713.
  27. Thiem U, Niklaus D, Sehlhoff B, et al. C-reactive protein, severity of pneumonia and mortality in elderly, hospitalised patients with community-acquired pneumonia. *Age Ageing* 2009;38:693–697.
  28. Kucukardali Y, Oncul O, Kunter E, et al. Community acquired infections in elderly population. *Cent Eur J Med* 2009;4:171–178.
  29. Lerner AJ, Hedera P, Koss E, Stuckey J, Friedland RP. Delirium in Alzheimer disease. *Alzheimer Dis Assoc Disord* 1997;11:16–20.
  30. Fong TG, Jones RN, Shi P, et al. Delirium accelerates cognitive decline in Alzheimer disease. *Neurology* 2009;72:1570–1575.
  31. van Gool WA, van de Beek D, Eikelenboom P. Systemic infection and delirium: when cytokines and acetylcholine collide. *Lancet* 2010;375:773–775.
  32. Aalten P, Verhey FR, Boziki M, et al. Neuropsychiatric syndromes in dementia: results from the European Alzheimer Disease Consortium: part I. *Dement Geriatr Cogn Disord* 2007;24:457–463.
  33. Hwang TJ, Masterman DL, Ortiz F, Fairbanks LA, Cummings JL. Mild cognitive impairment is associated with characteristic neuropsychiatric symptoms. *Alzheimer Dis Assoc Disord* 2004;18:17–21.
  34. Palmer K, Di Iulio F, Varsi AE, et al. Neuropsychiatric predictors of progression from amnesic-mild cognitive impairment to Alzheimer's disease: the role of depression and apathy. *J Alzheimers Dis* 2010;20:175–183.



## Neurology® Podcast

### Does vigorous exercise have a neuroprotective effect in Parkinson disease?

At [www.neurology.org](http://www.neurology.org), click on the “Download Latest Issue” link or “Subscribe Now” to subscribe to the RSS Feed.

This week's podcast features an interview by Dr. John Morgan with Dr. Eric Ahlskog about his paper on the effect of vigorous exercise on Parkinson disease.

**NEW CME Opportunity:** Listen to this week's Neurology® Podcast and earn 0.5 AMA PRA Category 1 CME Credits™ by answering the multiple-choice questions in the online Podcast quiz.



Scan this code with your smart phone to directly access the Podcast